

3485

~~SECRET~~

The Use of Aircraft in the Application of Chemicals to Agricultural
Crops in the Soviet Union

CIA HISTORICAL REVIEW PROGRAM
RELEASE AS SANITIZED
1998

I. Introduction

A. General

The Soviet Government currently is devoting considerable attention to the use of chemicals as a means of increasing agricultural production. The amount of mineral fertilizer allocated to Soviet agriculture is supposed to increase from 10.6 million metric tons in 1958 to 31 million metric tons by the end of the Seven Year Plan in 1965 (Selukoye Khomyaystvo, 15 March 1959, p. 3). Although currently the amount of herbicides and insecticides used in Soviet agriculture is relatively small, considerable attention is also being given to the combatting of agricultural pests and weeds by the use of chemicals. Information is not available concerning the amounts of insecticides and herbicides produced at present or concerning the planned production of these chemicals in 1965.

B. Volume of Work Done with Aircraft

The use of aircraft in the application of chemicals to agricultural crops in the Soviet Union has been increasing rapidly, having more than doubled during the past five years. The following data on the volume of aviation-chemical work show the increasing importance of aircraft as a means of applying chemicals to agricultural crops in the USSR:

~~SECRET~~

<u>Year</u>	<u>Increase over 1940^{1/}</u> <u>(percent)</u>	<u>Area covered^{2/}</u> <u>(1,000 hectares)</u>
1940	100	1,000
1950	350	3,500
1951	400	4,000
1952	500	5,000
1953	600	6,000 ^{2/}
1954	700	7,000 ^{2/}
1955		11,000 ^{3/}
1956		16,200 ^{4/}
1960 (plan)		26,500 ^{3/}
1965 (plan)		32,500 ^{4/}

The Soviets plan to double the area of agricultural crops on which chemicals are applied from airplanes between 1958 and 1965. Although the Soviets have generally fallen far short of past planned goals in agriculture and related fields, the use of aircraft in the application of chemicals to agricultural crops in the Soviet Union is expected to continue to increase rapidly in the years ahead. The generally larger size of fields and farming units in the Soviet Union than in the United States and other countries tend to make aircraft a relatively more economical method of applying chemicals to agricultural crops than in the other countries.

C. Use of Aircraft in Soviet Agriculture

The different types of work in the agricultural field being done by aircraft in the Soviet Union are numerous. According to the Soviets, even in 1954, aviation-chemical work at least in Ukrainian agriculture already was ceasing to have a seasonal character.^{5/} Also, they state that in Uzbekistan "air-chemical work is now carried out the year round, whereas until 1954 it

was of a seasonal nature".^{6/} Among the more important operations in which aircraft are used in applying chemicals to agricultural crops are the application of mineral fertilizers to fall sown (winter) grains, primarily winter wheat, as well as to a number of other crops such as cotton, and sugar beets; the application of herbicides to crops such as small grains and flax; the application of insecticides to a number of different crops and to forests; and the application of defoliants to cotton. Other types of work carried out by aircraft in the general field of agriculture include the spraying of vineyards and orchards, the seeding of forest and desert areas, patrol flights protecting forests areas from fire, extermination of wolf packs in the cattle raising areas of Siberia, and hunting fur-bearing animals in the Far East.

D. Published Information on Movement of Agricultural Aircraft

Members of the U.S. Cotton Growing Delegation who toured the USSR in 1958 were told that the aircraft used in Soviet agriculture were centrally controlled and dispatched. Also, it is stated that "in contrast to the prewar period, - - - - a schedule of operations was set up which followed the course of agricultural work throughout the various geographical areas."^{7/} Little information is published, however, on the actual movement of aircraft equipped for applying chemicals between the different agricultural areas of the Soviet Union. An article on pages 6-7 of the no. 3, 1958 issue of the journal Civilian Aviation contained the statement that "Besides the

high-quality work conducted in the Ukraine, the same flight commanded by Babich dispatches airplanes for arial treatment of cotton fields in Uzbekistan and Kirgiziya." A broadcast from Kiev stated that "a group of AN-2 planes of the Ukrainian Civil Aviation Fleet equipped with fertilizer spreading devices left for Uzbekistan on May 29 to help the cotton growers. They will be working in Andizhan, Namangan, and (Surhant?) Oblasts. It is planned to send such groups also to Krasnodar Krai and Azerbaydhan." 8/ Another broadcast, this one from Leningrad, stated that "Ten AN-2 aircraft with equipment for combating agricultural pests will be sent by the Northern Administration of Civil Aviation to the Virgin Lands of Kazakhstan. ---- Preparations for the flight from Leningrad to Kazakhstan (Kustanal and Pavlodar Oblasts mentioned) began on April 6 at Leningrad Airport." 2/ Finally, a Pavlodar broadcast on April 24, 1959 stated that "Every year the Northern Directorate of the Civil Aviation Fleet sends airplanes to the Virgin Lands to help with weed and pest killings.----- the planes will bring as May Day gifts books and musical instruments for field mechanizers." 10/

B. Limitations of the Study

The research work carried out on this project has resulted in a number of limitations of the study becoming apparent. Very little information on the dates at which the various crops reach the different phenological stages in the principal agricultural areas of the USSR was found in Soviet

literature. The dates at which the various crops are normally planted in the different regions could not be found in the basic literature on Soviet Agriculture. Thus, the data upon which this study is based represent one of the more pronounced limitations of the study.

Phenological data on winter rye, winter and spring wheat, and spring barley were obtained from the following books by M.Y. Mattonson of the American Institute of Crop Ecology: Rye - Climate Relationships and the Use of Phenology in Ascertaining the Thermal and Photo-Thermal Requirements of Rye, Washington, D.C., 1953; Wheat-Climate Relationships and the Use of Phenology in Ascertaining the Thermal and Photo-Thermal Requirements of Wheat, Washington, D.C., 1955; and Barley - Climate Relationships and the Use of Phenology in Ascertaining the Thermal and Photo-Thermal Requirements of Barley, Washington, D.C., 1957. These studies are based upon data which were obtained from agricultural experiment stations in the Soviet Union and pertain to a period some 30-40 years ago. The question immediately arises, does the experiment station data accurately reflect the phenological stages for crops in the area or could the differences in agricultural practices and in the seed varieties used on the experiment stations as opposed to those in general use in the agriculture of the area have caused a significant variation in the phenological stages of the crops? Also, have technological changes and the development of new seed varieties during the past several decades caused a significant change in the duration of the various phenological

stages of the different crops? Answers to these questions could not be found in recent literature published by the Soviets.

Information on the approximate sowing and harvesting dates for oats was obtained from an ORR report published in June 1952. Additional information on this report can be obtained, if needed, from ORR/R/AG. The table used from this report as well as the appropriate pages from the Suttonson studies are contained in Appendix C. Reference to maps from NIS 26, Section 61, "Agriculture, Fisheries, and Forestry," April 1958, Figures 61-10, 11, 14, 15, 17, 22, 62, and 68, showing acreage distribution of various crops in the USSR, may be useful to readers of this report.

The lack of phenological information also pertained to the technical crops considered in this study. In order to estimate the ranges of probable planting dates for these crops, it was necessary to study spring planting progress reports in the Soviet press for the past 3-4 years.

~~SECRET~~

Despite a lack of basic information the conclusions reached in this study will serve as a useful guide in determining when aircraft are most likely to be used in applying chemicals to agricultural crops in the Soviet Union.

~~SECRET~~

II. Methodology

A. General

Phenological data on several of the small grains (wheat, rye, and barley) were obtained from the above mentioned studies by H.Y. Ruttonson. Ruttonson's studies are based on data from North America and thermally analogous areas in the Soviet Union and other European countries. Average and extreme dates for the sowing, heading, and ripening of winter rye, and winter and spring wheat, and spring barley in the principal agricultural areas of the USSR were determined from these data. This was done by inspecting the data given for individual experiment stations within each agricultural area in the USSR. In some instances, where insufficient data were available, it was necessary to consider data from stations outside but immediately adjacent to the agricultural area in question.

Phenological data for oats were not available as such and had to be derived from planting and harvesting information in the above mentioned ORR report. Since all the small grains have similar phenologies, that for oats was calculated by applying general phenological information, provided by a US crop specialist at Beltsville, to the planting and harvesting dates.

In addition to general information concerning the phenology of the

various crops, specialists at Beltsville also provided information on the stages of development during which the various types of agricultural chemicals can or would most likely be applied to these crops under normal conditions.

As indicated above, phenological data were unavailable from Soviet publications for the technical crops (fiber flax, sugar beets, sunflowers, and cotton) considered in this study. In order to determine approximate dates for the different stages of development for these crops, it was necessary to (1) scrutinize Soviet press reports on spring seeding progress for the past 3-4 years in order to determine the range of dates during which each of the technical crops is normally planted; (2) apply the general phenological information supplied by the Beltsville specialists to these planting periods. In this manner, approximate dates for the various phenological stages pertinent to the application of agricultural chemicals were obtained.

B. Agricultural chemicals

To achieve maximum effectiveness and to avoid possible deleterious effects on crop growth and yields, agricultural chemicals should be applied only when the plants are in certain stages of development. The times at which these chemicals are best applied vary not only according to whether a crop is of the narrow-leaf (small grains and fiber flax) or broad-leaf (sugar beets, sunflowers, and cotton) type but also according to type of chemical (herbicides, insecticides, mineral fertilizer, or defoliant) applied.

1. Herbicides

Herbicides achieve their greatest utility in controlling weeds in crops which are members of the grass family such as grains and fiber flax. The grain crops are not particularly susceptible to herbicidal damage except during the critical tillering, blooming, and heading stages of development.*

* The application of herbicides to corn has not been considered in this study. Phenological data are not available for corn in the USSR, and it is believed that herbicides would not generally be applied to corn by airplanes.

To be most effective in controlling weeds and yet not cause yields to be adversely affected, herbicides should be applied to small grains during the jointing stage. However, no direct information is available on the dates of the jointing stages. Heading and ripening dates for the small grains in the different regions of the USSR were used as points of reference in calculating the ranges of dates during which herbicidal treatment would most likely take place. The heading and ripening were taken from Buttomson and are given in Appendix C. It was assumed that jointing starts 7 weeks prior to ripening and lasts 3 weeks, for all small grains and in all areas.

The approximate phenological dates given in Appendix B were used in calculating the range of possible dates during which herbicidal spraying would be carried out on the small grains. It was assumed that the beginning of spraying coincides with the onset of jointing and continues until one week before heading.

The similarity of the phenologies of the different small grains sown in the spring and of the winter grains after the tillering stage resulted in the use of the same criteria for all these crops in determining the ranges of dates during which herbicides would most likely be applied. Differences in phenologies between the individual grains are generally less than those which occur between varieties of the same grain. Varietal differences, according to the Beltsville specialists, can cause up to two weeks difference either way in the phenological development of a grain.

Fiber flax differs from the small grains somewhat in that it is usually treated with herbicides at a much earlier stage of development. As opposed to the small grains, fiber flax is most effectively sprayed with herbicides when it is 2-6 inches tall. It attains this height 3-4 weeks after emergence, or about 4-6 weeks after planting. This is the criteria employed in setting the range of dates when herbicides would most likely be applied to fiber flax.

Sugar beets, sunflowers, and cotton, all broad-leaf crops, are generally not treated with herbicides during the growing season. These crops are highly susceptible to damage by herbicides. If these crops are to be sown in a particularly weedy area and chemical weed control is desired, the common practice is to apply herbicides prior to the time of emergence. (Sugar beets, sunflowers, and cotton emerge 1-2 weeks after planting). The only available

information contrary to this indicates that cotton in the USSR is sometimes sprayed with an herbicide in conjunction with cultivation, implying the chemical would be applied from the ground. No information whatever is available to suggest that aircraft are used during the growing season to apply herbicides to sugar beets, sunflowers, or cotton. Thus, it has been assumed that aircraft are not used to apply herbicides to these crops in the Soviet Union.

2. Insecticides

Insecticides are applied to a relatively wide range of crops in the Soviet Union. They generally can be applied at almost any stage of crop development, depending upon time of insect infestation, without harmful effects. Aircraft are becoming more widely used in applying insecticides as well as the other agricultural chemicals. According to the Soviets, the use of aircraft as a means of application results in more efficient utilization of available agricultural chemicals which are in relatively short supply in the USSR.

With respect to the small grains, the ranges of dates when insecticides could be applied, found in Appendix A, were based on the premise that if it were necessary to spray a crop, it could be done anytime between emergence (about 1-2 weeks after sowing) and about a week or so prior to ripening. Spraying later than this might possibly result in a toxic residue being left on the harvested grain.

Fiber flax, according to a specialist at Beltsville, rarely suffers a severe insect problem. If spraying is required, however, it could be carried out during the same stages of development as for the small grains.

In the case of sugar beets, which are frequently and severely infested with beet weevils in the Ukraine, insecticides can probably be effectively applied from aircraft anytime between emergence and the time the leaves are sufficiently large to cover the soil surface, approximately 3-3 1/2 months after planting. If an infestation were to occur subsequent to this stage of development, eradication would probably be most effective if insecticides could be applied to both the top and bottom leaf surfaces. Thus, it is assumed that application of insecticides after the plants are 3-3 1/2 months old would be made by ground equipment.

Sunflowers are most susceptible to insect attack when the plant is about two months old. Flower head formation is beginning about that time, and it has been assumed that insecticides would be applied then.

A cotton specialist at Beltsville who visited the Soviet Central Asian cotton growing republics in 1956, stated that if the Soviet cotton crop becomes infested, the most likely time to apply insecticides is from early July to early September. This is the criteria that has been used in setting the ranges of dates found in Appendix A.

3. Mineral Fertilizers

In the past, the technical crops have been the principal

claimants of the limited amounts of mineral fertilizers available in the USSR. However, fertilization of winter grain crops has been increasing in recent years. Spring crops other than the technical crops also receive applications of mineral fertilizer. Recent increases in the supply of mineral fertilizer is resulting in a wider variety of crops being fertilized. Most of the mineral fertilizers applied to spring crops, including spring-sown grain, technical crops, potatoes, and vegetables, are applied at the time of soil preparation or planting. Subsequent side-dressings of mineral fertilizers, in general, are applied by a surface means to row crops. Aircraft are used in applying mineral fertilizers mainly to winter grain and certain technical crops.

Applications of mineral fertilizers from aircraft to winter grain, primarily winter wheat,* are made both in the fall and spring. It is

* It is believed that most of the mineral fertilizers applied to winter grains is applied to winter wheat because the returns are greater.

believed, however, that most of this work is performed in the spring.

With respect to determining the times when mineral fertilizers would most likely be applied from aircraft to winter wheat, the following criteria were used. In order for winter wheat to benefit from mineral fertilizers applied during the fall months, application should be made well in advance of the end of the growing season. It has been assumed,

therefore, that this type of agricultural chemical would be applied by aircraft over a four week period centered around a date about six weeks before the snow cover normally becomes established in each of the principal winter wheat regions of the USSR. Also, the assumption has been made that mineral fertilizers applied to winter wheat in the spring would most likely be put on over a four week period centered around a date about two weeks after the snow cover disappears.

The dates when the snow cover normally becomes established and disappears are important in determining the most likely times when mineral fertilizers would be applied to winter grain. Average dates for these natural phenomena for the principal winter wheat regions (the Ukraine, North Caucasus, and Central Black Soil Zone) were estimated from data in MIS 26, Section 23, "Weather and Climate of the USSR."

Little evidence was found that aircraft are used in applying mineral fertilizers to spring sown crops in the Soviet Union. In 1952 and 1953 aircraft reportedly were used in applying mineral fertilizers to relatively small acreages of cotton and sugar beets. 11/ No evidence was found that aircraft are used in applying mineral fertilizers to spring sown grains or fiber flax. However, if mineral deficiencies become evident during the growing season in these crops, aircraft would probably be the best means for applying fertilizers. For the purpose of this study, it has been assumed that, if aircraft are used in applying mineral fertilizers to spring sown

crops, their use for this purpose would roughly correspond to the periods when insecticides would most likely be applied, as shown in Table 1.

4. Defoliants

The use of aircraft in applying defoliants to cotton in the USSR has been increasing rapidly in recent years. The Soviets reported that in 1951 about 80% of the cotton in Uzbekistan, where about two-thirds of Soviet cotton is produced, was defoliated by aerial spraying. 12/ Currently, the Soviets state that ground equipment should be used for spraying defoliants on cotton only in those areas inaccessible to aircraft by reason of trees, power lines, etc. 13/ Thus, it may be assumed that a large part of the cotton in the Soviet Union is defoliated by means of aerial spraying.

Defoliants are sprayed on the cotton crop at the time the bolls begin to open. In the Soviet Union, this normally occurs from about mid-September until about mid-October. This has been verified by the cotton specialist from Beltsville who visited the main USSR cotton growing areas in 1953 and witnessed the beginning of aerial application of defoliants in mid-September.

III. Seasonal and Geographic Pattern of Aerial Spraying and Dusting.

A summary of the periods during which various types of agricultural chemicals are most likely to be applied to selected crops in the principal agricultural areas of the Soviet Union, as given in Appendix A, shows that this type of work can be carried out over extended periods. The application of chemicals to agricultural crops is confined to the growing season. Thus, little, if any, of this work is performed during the period from about November 1 to March 1. Also, this summary shows that in some regions, namely the Ukraine, North Caucasus, Volga, Urals, and Northern Kazakhstan, a period covering on an average the first half of September would be relatively free from work in the application of chemicals to agricultural crops.

This summary also reveals that the periods during which the various types of agricultural chemicals could be applied to crops in the Soviet Union are roughly as follows: mineral fertilizers would be applied to winter wheat in the spring from about mid-March to mid-May with most of the work being done in April; insecticides could be applied during a six month period from early March until early September; herbicides could be applied from about mid-April until mid-August; defoliants would most likely be applied to the cotton between about mid-September and mid-October; and the first application of mineral fertilizers would be applied to winter wheat in the fall, between the early part of September and the latter half of October.

Although these agricultural chemicals can be applied to the various crops over rather extended periods, there are some factors which are believed to impose limitations on the periods during which aircraft would most likely be used extensively in applying the chemicals. First of all, the supply of agricultural chemicals in the Soviet Union is limited so that in general only a part of the total acreage devoted to a crop can be treated. For example, only about one percent of the grain acreage in 1957 was treated with herbicides.^{1b/} Secondly, the largest part of a given crop in a particular area would reach the appropriate stage of development for applying chemicals over a shorter range of time than indicated by the extreme ranges in dates as given in Appendix A. Only a relatively small part of the crop is believed to reach the appropriate stage for treatment toward either end of the extreme ranges in dates when chemicals can be applied. Thus, the middle third of the extreme ranges is felt to be the period when most of the work in applying agricultural chemicals would be carried out.

It has, therefore, been assumed that the extensive use of aircraft in applying agricultural chemicals in general is limited to the middle third of the extreme ranges in dates when chemicals can be applied to crops in a particular area. Those shorter periods during which aircraft are most likely to be used extensively in applying herbicides and insecticides as well as mineral fertilizers to crops in the principal agricultural areas of the Soviet Union are presented in Table 1.

Table 1. Periods During Which Aircraft are Most Likely to be Used Extensively in Applying Chemicals to Crops in the Principal Agricultural Areas of the Soviet Union

	Herbicides	Insecticides	Mineral Fertilizers	
			Fall	Spring
Belorussia	5 Jun 15 Jul	5 Jun 1 Aug		
Central Non-Black Soil Zone	25 May 30 Jun	25 May 15 Jul		
Central Black Soil Zone	25 May 25 Jun	25 May 15 Jul	1 Sep 1 Oct	1 Apr 1 May
Ukraine and Moldavia	15 May 20 Jun	12 May 10 Jul	15 Sep 15 Oct	1 Apr 1 May
North Caucasus	20 May 15 Jun	10 May 25 Jun	1 Oct 1 Nov	15 Mar 15 Apr
Volga	1 Jun 5 Jul	1 Jun 25 Jul		
Urals	5 Jun 5 Jul	1 Jun 20 Jul		
West Siberia	10 Jun 20 Jul	5 Jun 25 Jul		
Kamchatka (Northern Part)	15 Jun 15 Jul	1 Jun 20 Jul		
Central Asia	5 May 5 Jun	5 May 1 Sep	10 Sep	15 Oct

Defoliation

~~SECRET~~

Sources:

- a/ Further information on the procedures used in this research project is contained in the Methodology section of this report.
- b/ These periods for applying chemicals are believed applicable to the southern part of Kazakhstan as well as to the Uzbek, Kirgiz, Tadzhik, and Turkmen Republics.

~~SECRET~~

~~SECRET~~

For convenience in discussing the seasonal and geographic pattern of aerial spraying and dusting the ten agricultural areas under consideration have been combined into four groups.*

*It should be noted that the Transcaucasus has not been included among the agricultural regions considered in this study. Even though it is adjacent to several of the regions considered it was not included because, in relation to Soviet Central Asia, the Transcaucasus is of minor agricultural importance except for such crops as tea and citrus fruits. Inasmuch as the periods during which aircraft might be used in applying chemicals to crops in the Transcaucasus would roughly coincide with the most intensive requirement in Central Asia, it is believed that there would be little likelihood of agricultural aircraft being transferred between these two regions.

Belorussia, the Central Non-Black Soil Zone, the Volga Region, and the Urals comprise the first group. The extensive use of aircraft in applying herbicides and insecticides to crops in these areas would be largely confined to a period extending from about 1 June to the second half of July. Aircraft reportedly were used to apply herbicides to fiber flax in the Central Non-Black Soil Zone during the 1958 harvesting season.¹⁵ Normally haying takes place in this area around the latter half of June. As indicated in Table 1, herbicides would most likely be applied from aircraft between 25 May and 30 June in the Central Non-Black Soil Zone.

The second group is made up of the Ukraine, the Central Black Soil Zone, and the North Caucasus. Because these regions are the principal winter wheat

~~SECRET~~

growing areas of the USSR, it is believed that the extensive use of aircraft in applying mineral fertilizers to winter grains is largely confined to these areas. Aerial fertilization in the spring is likely to be most extensive from about mid-March through April and in the fall during September and October. However, it has been reported that aircraft have been used for this purpose in the southern Ukraine as early as February and in the fall as late as November and December.^{16/} Also, in the March 1953 issue of *Grachanskaya Aviatsiya* (Civilian Aviation) pages 6-7, it was reported that aerial fertilization work had already started in Kiev and Cherkassy Oblasts.

In the North Caucasus, fertilization of winter crops on the Gigant state farm started on 10 March in 1947.^{17/} Also, by 1 April 1953, almost 70,000 hectares of crops, primarily winter wheat, (of a total of 215,000 hectares planned to be fertilized during the year) had been fertilized from the air by the Stavropol Detachment of Agricultural Aviation.^{18/}

In the Ukraine, the Central Black Soil Zone, and the North Caucasus aerial spraying of insecticides and herbicides is believed largely confined to the period extending from about mid-May to mid-July. Thus, it is believed that the extensive use of aircraft in the spring application of mineral fertilizers and in the application of herbicides and insecticides in these regions would be completed by about mid-July and that many of the aircraft used during these periods of peak activity could be dispatched to other areas.

The third area for discussion includes West Siberia and the Northern part of Kazakhstan. Aerial treatment of crops in this area probably is confined largely to the spraying of insecticides and herbicides. Extensive use of aircraft for this work probably would be confined largely to a period extending from about June 1 until the latter part of July. Experimental aerial spraying of herbicides on (spring) wheat was reportedly conducted on two state farms in Kustanaisky and Karabalytsky Raions of Eastern Oblast during the second half of June in 1958.^{39/} Also, a radio broadcast from Alma Ata on 9 April 1959 stated that both airplanes and tractors were to be used to apply insecticides during the spring field work campaign in combatting grain stem borers in the main virgin land regions of Kazakhstan.^{29/} As indicated earlier in the Published Information on Movement of Agricultural Aircraft section of this report, aircraft equipped for applying agricultural chemicals were sent from Leningrad to Northern Kazakhstan during the latter part of April.

The fourth area to be discussed is Soviet Central Asia, including the Uzbek, Kirgiz, Tadzhik, and Turkmen Republics and the southern part of Kazakhstan. The agricultural economy of this area is centered around cotton growing and it is in the treatment of this crop that most of agro-chemical work is performed. However, there are other crops in the area on which herbicides and insecticides are normally required. The use of aircraft in the application of chemicals to the crops other than cotton is believed to be of relatively

less importance and confined largely to May and June.

Aircraft are used extensively to apply insecticides to the cotton fields from about mid-July through August. According to the Beltsville cotton specialist, insects appear to be well controlled in Soviet cotton fields. He was told that some cotton fields in Azerbaydzhan had received as many as 10 applications of DDT between mid-July and 1 September. However, on 10 June 1959 it was reported from Tashkent that AN-2 planes and pilots sent by the Ukrainian administration of the Civil Air Fleet had immediately begun spraying the cotton fields after arriving in Namangan Gulast.²¹ Probably this early transfer of aircraft from the Ukraine to Soviet Central Asia can be attributed to the early spring in the Soviet Union in 1959.

As indicated in the ~~Entomology~~ ^{Phytology} section of this report, aircraft are not used in applying herbicides to cotton. The period from about mid-September to mid-October is one of intense aerial activity when the cotton fields are being treated with defoliants.

Sources:

- 1/ "Civilian Aviation in Agriculture", Moscow, 1958, p. 6.
- 2/ *Grazhdanskaya Aviatsiya* (Civilian Aviation), No. 31, November 1958, p. 5.
- 3/ Air Intelligence Information Report, IR-1643-57, 6 Aug 57, p. 2.
- 4/ Air Intelligence Information Report No. 1235578, 31 July 1959, p. 2.
- 5/ *Civilian Aviation*, op. cit., p. 43.
- 6/ *Grazhdanskaya Aviatsiya* (Civilian Aviation), No. 5, 1956, p. 32.
- 7/ *Zashchita Rasteniy Ot Vreditel'ov i Bolesnei* (The Protection of Plants Against Pest and Diseases), No. 6, Moscow, Nov/Dec., p. 39.
- 8/ FBIS 59, M3243, ORO.
- 9/ FBIS 59, L3876, ORO.
- 10/ FBIS 59, L4406, ORO.
- 11/ "Civilian Aviation in Agriculture," op. cit., pp. 28-30.
- 12/ Ibid., p. 68.
- 13/ *Khlopkovodstvo* (Cotton Growing), No. 9, 1959, p. 3.
- 14/ *Zashchita Rastenii ot Vreditel'ei i Bolesnei* (Protection of Plants from Pests and Diseases), No. 5, Sept. - Oct., 1959, p. 18.
- 15/ *Nauka i Zhizn* (Science and Life), No. 3, 1959, p. 30.
- 16/ "Civilian Aviation in Agriculture," op. cit., p. 43.
- 17/ Ibid., p. 51.
- 18/ *Selskoye Khozyaystvo*, 5 April 1959, p. 4.
- 19/ *Zemledeliye*, no. 5, 1959, pp. 36-39.
- 20/ FBIS 59 L 3970, 23 April 1959, ORO.
- 21/ CIA, FID Summary No. 2260, 19 Aug 1959, pp. 2-3, ORO.

Appendix A. Periods During which Various Types of Agricultural Chemicals May be Applied to Selected Crops in the Principal Agricultural Areas of the Soviet Union.^{8/}

Crop	Herbicides		Insecticides		Mineral Fertilizer	
					Fall	Spring
<u>Moldavia</u>						
Winter Rye	25 Apr	15 Jun	20 Apr	10 Aug	15 Sep	15 Oct 1 Apr 1 May
Oats	20 Jun	25 Aug	15 Apr	25 Sep		
Fiber Flax	25 May	20 Jul	15 May	15 Aug		
<u>Central Non-black Soil Zone</u>						
Winter Rye	20 Apr	10 Jun	5 Apr	5 Aug	1 Sep	1 Oct 15 Apr 15 May
Spring Barley	5 Jun	10 Jul	10 May	10 Aug		
Oats	25 Jun	1 Aug	25 Apr	2 Sep		
Fiber Flax	25 May	20 Jul	15 May	15 Aug		
<u>Central Black Soil Zone</u>						
Winter Rye	25 Apr	30 May	20 Apr	10 Aug	1 Sep	1 Oct 1 Apr 1 May
Winter Wheat	10 May	20 Jun	25 Apr	25 Jul	1 Sep	1 Oct 1 Apr 1 May
Spring Wheat	25 May	20 Jun	10 Apr	15 Aug		
Spring Barley	5 May	1 Jul	5 Apr	10 Aug		

(continued)

Appendix A. Periods During Which Various Types of Agricultural Chemicals May Be Applied to Selected Crops in the Principal Agricultural Areas of the Soviet Union. ^{2/} (continued)

Crop	Herbicides		Insecticides		Mineral Fertilizer	
					Fall	Spring
<u>Central Black Soil Zone</u>						
Oats	25 Jan	1 Aug	15 Apr	1 Sep		
Sugar Beets			15 Apr	1 Sep		
Sunflowers			1 Jan	15 Aug		
Urticaria						
Winter Rye	10 Apr	30 May	25 Mar	25 Jul	15 Sep	15 Oct
Winter Wheat	25 Apr	20 Jun	1 Apr	1 Aug	15 Sep	15 Oct
Spring Barley	5 May	15 Jun	20 Mar	10 Aug		
Oats	20 May	1 Jul	20 Mar	1 Aug		
Fiber Flax	25 May	20 Jul	15 May	15 Aug		
Sugar Beets			15 Apr	1 Sep		
Sunflowers			1 Jan	15 Aug		

(continued)

Appendix A. Periods During which Various Types of Agricultural Chemicals May be Applied to Selected Crops in the Principal Agricultural Areas of the Soviet Union. 2/ (continued)

Crop	Restrictions			Mineral Fertilizers		
	Restrictions			Fall	Spring	Spring
<u>North Caucasus</u>						
Winter Wheat	30 Apr	30 Jun	20 Apr	1 Aug	1 Oct	1 Nov
Spring Wheat	15 May	15 Jun	5 Apr	25 Jul		15 Mar
Spring Barley	25 May	5 Jul	1 Apr	1 Aug		15 Apr
Oats	20 May	20 Jun	25 Mar	20 Jul		
Sunflowers			1 Jun	15 Aug		
<u>Volga</u>						
Winter Rye	30 Apr	5 Jun	20 Apr	20 Jul	15 Sep	15 Oct
Winter Wheat	5 May	15 Jun	5 Apr	15 Jul	1 Apr	1 May
Spring Wheat	10 May	10 Jul	15 Apr	1 Sep	15 Sep	1 May
Spring Barley	30 May	20 Jun	20 Apr	20 Jul		
Oats	15 Jun	20 Aug	15 Apr	20 Sep		
Sunflowers			1 Jun	15 Aug		

(continued)

Appendix A. Periods during which Various Types of Agricultural Chemicals May be Applied to Selected
Crops in the Principal Agricultural Areas of the Soviet Union. 2/ (continued)

Crop	<u>Pesticides</u>		<u>Insecticides</u>		<u>Mineral Fertilizer</u>	
					<u>Fall</u>	<u>Spring</u>
<u>Grains</u>						
Winter Rye	5 May	20 Jun	10 Apr	1 Aug	15 Sep	15 Oct 15 Apr 15 May
Spring Wheat	25 May	5 Jul	15 Apr	5 Sep		
Oats	25 Jun	5 Aug	15 Apr	5 Sep		
Fiber Flax	25 May	20 Jul	15 May	15 Aug		
Scarifiers			1 Jun	15 Aug		
<u>East Siberia</u>						
Winter Rye	15 May	20 Jun	1 May	1 Aug	15 Aug	15 Sep 1 May 1 Jun
Spring Wheat	15 Jun	25 Jul	5 May	10 Sep		
Spring Barley	25 May	10 Jul	15 May	15 Aug		
Oats	25 Jun	5 Aug	15 Apr	5 Sep		
Fiber Flax	25 May	20 Jul	15 May	15 Aug		

(continued)

Appendix A. Periods During which Various Types of Agricultural Chemicals May be Applied to Selected Crops in the Principal Agricultural Areas of the Soviet Union.^{b/} (continued)

Crop	Pesticides		Fertilizers	
			Mineral Fertilizers	Organic Fertilizers
			Fall	Spring
<u>Russia</u>				
Winter Rye	15 May	5 Jul	10 Apr	1 Aug
Spring Wheat	25 May	10 Jul	25 Apr	26 Aug
Spring Barley	30 May	20 Jun	10 Apr	15 Aug
Oats	1 Jul	(10 Aug)	15 Apr	(1 Sep)
Sunflowers			1 Jun	15 Aug
<u>Soviet Central Asia^{b/}</u>				
Winter Wheat	5 Apr	25 May	5 Mar	1 Jul
Spring Wheat	30 Apr	5 Jun	25 Feb	5 Jul
Oats	5 Jun	(5 Jul)	5 Apr	(15 Aug)
Cotton	10 Sep ^{c/}	15 Oct ^{c/}	1 Jul	1 Sep

1-2-3-4-5-6-7-8-9-10-11-12-13-14-15-16-17-18-19-20-21-22-23-24-25-26-27-28-29-30-31-32-33-34-35-36-37-38-39-40-41-42-43-44-45-46-47-48-49-50-51-52-53-54-55-56-57-58-59-60-61-62-63-64-65-66-67-68-69-70-71-72-73-74-75-76-77-78-79-80-81-82-83-84-85-86-87-88-89-90-91-92-93-94-95-96-97-98-99-100

1-2-3-4-5-6-7-8-9-10-11-12-13-14-15-16-17-18-19-20-21-22-23-24-25-26-27-28-29-30-31-32-33-34-35-36-37-38-39-40-41-42-43-44-45-46-47-48-49-50-51-52-53-54-55-56-57-58-59-60-61-62-63-64-65-66-67-68-69-70-71-72-73-74-75-76-77-78-79-80-81-82-83-84-85-86-87-88-89-90-91-92-93-94-95-96-97-98-99-100

Footnotes to Appendix A.

- a/ The periods for applying herbicides and insecticides to the various crops were derived by relating information concerning the best or most effective time (stage of plant development) of application obtain from various crop specialists at the U.S. Agricultural Experiment Station at Beltsville to the dates at which the crops in question reach the different phenological stages as contained in Appendix B. Further details on the procedure used in establishing the periods for applying herbicides and insecticides to the various crops, as well as that used in estimating the periods when mineral fertilizers would be applied by aircraft, are contained in the methodology section to this report.
- b/ Includes the Republics of Uzbekistan, Kirgiz, Tadzhik, and Turkmen. Due to proximity and the similarity of cropping patterns, the probable dates for applying chemicals in Soviet Central Asia would also apply to southern Kazakhstan.
- c/ Period during which defoliant is applied to cotton.

Note

Dates in parenthesis are estimates based upon approximate duration of the periods in other regions. Ending dates for the harvesting season were not available for oats in the regions concerned.

Appendix B. Average in 1956 and Available Phenological Dates (Approximate) for Selected Crops in the
 Potential Agricultural Regions of the Soviet Union² (continued)

Crop	1956 Average		Date Sown		Date Harvested		Harvest Date	
	(1,000 ha.)	Percent of Under Total	Average	Extremes	Average	Extremes	Average	Extremes
Wheat								
Winter Rye	2,231	10.1			5 Jun	20 May 20 Jun	27 Jul	3 Jul 9 Aug
Spring Wheat	6,517	19.3	5 May	10 Apr 20 May	27 Jun	14 Jun 20 Jul	2 Aug	13 Jul 20 Sep
Oats	1,769	11.6	22	10 Apr 10 Jun	22	22	22	15 Aug 20 Sep
Winter Flax	21	4.9						
Buckwheat	272	6.0	22	20 May 1 Jun				
Total 1956 From Area	17,191	31.0						
Barley								
Winter Rye	716	3.9			27 Jun	6 Jun 20 Jun	20 Jul	21 Jul 9 Aug
Spring Wheat	10,613	21.6	24 May	20 Apr 10 Jun	15 Jul	6 Jul 20 Jul	22 Aug	9 Aug 1 Oct 9/
Grain Barley	432	3.9	20 May	10 May 20 Jun	17 Jul	12 Jul 20 Jul	6 Aug	20 Jul 22 Aug
Oats	2,889	19.0	22	10 Apr 20 Jun	22	22	22	15 Aug 20 Sep
Winter Flax	205	5.5	22	20 Apr 15 Jun				
Total 1956 From Area	20,709	10.6						

(continued)

Footnotes to Appendix 2.

- 1/ Average data for 1976 were obtained from various pages in Volume I and II of the Soviet statistical handbook, *Posernyye Vloshchadi SSSR* (Sown Area of the USSR), Moscow, 1977. The data upon which the phenological dates for the various grades are based will comprise Appendix 2 to this report and will be forwarded (type remaining 5 lines of footnotes as they appear in the original version).
- 2/ 1976 data.
- 3/ Data for the station at Smolensk, USSR, accounts for the 1 October date. Data for other stations suggest that very little spring wheat will ripen before September 13.
- 4/ Includes the Republics of Uzbek, Kirgiz, Tadzhik, and Turkmen.
- 5/ Phenological data available from Ashkhabad, Russian SSR, only.

Appendix B. Average in 1956 and Available Handedness Dates (Approximate) for Selected Crops in the Principal Agricultural Regions of the United States 5/ (continued)

Crop	1956 Average (1,000 ha.)	Percent of Total	Date Ending		Date Ending	
			Average	Minimum	Average	Minimum
Wheat						
Winter wheat	145	2.6			20 Jul	3 Jul 9 Aug
Spring wheat	17,954	33.0			15 Aug	14 Jul 8 Sep
Spring barley	2,387	4.0			1 Aug	21 Jul 23 Aug
Oats	720	1.2				20 Aug 22
Barley	320	0.7				
Total 1956 Farm Area	27,823	14.3				
Barley, Oats, and Rye						
Winter wheat	204	0.3			20 Jul	31 May 6 Jul
Spring wheat 5/	633	1.3			10 Jul	18 Jun 11 Jul
Spring barley 5/	857	2.4			20 Aug	22 22
Oats	59	0.4			22	25 Jul 22
Cotton	1,732	89.9			22	22 22
Total 1956 Farm Area	3,426	2.9				

Footnotes to Appendix B.

- a/ Average data for 1975 were obtained from various pages in Volumes I and II of the Soviet statistical handbook, *Posvayye Ploshchadi SSSR* (Basic Area of the USSR), Moscow, 1977. The data upon which the phenological dates for the various grains are based will appear in Appendix C to this report and will be forwarded (type retaining 5 lines of footnotes as they appear in the original version).
- b/ 1975 data.
- c/ Data for the station at Shadrinsk, USSR, accounts for the 1 October date. Data for other stations suggest that very little spring wheat will ripen after September 15.
- d/ Includes the Republics of Uzbek, Kirgiz, Tadzhik, and Turkmen.
- e/ Phenological data available from Ashkhabad, Turkmen SSR, only.

Appendix B. Average in 1936 and Available Phenological Dates (Approximate) for Selected Crops in the Principal Agricultural Regions of the United States

Crop	1936 Average (1,000 ha.)	Percent of 1936 Total	Date First		Date Second		Date Third	
			Average	Midwestern	Average	Midwestern	Average	Midwestern
<u>Midwestern</u>								
Winter Rye	1,469	8.0			2 Jan	10 May	22 Jan	25 Jul 12 Jul 15 Aug
Oats	576	3.0	11.2	10 Apr 30 May	11.2	11.2	11.2	11.2 20 Aug 30 Sep
Winter Wheat	340	17.7	11.2	25 Apr 13 Jun	11.2	11.2	11.2	
Total 1936 Area Acres	5,470	2.8						
<u>Central Non-Irrigated Soil Zone</u>								
Winter Rye	4,356	23.7			10 May	11 May	15 Jan	24 Jul 20 Jan 14 Aug
Spring Barley	262	2.3	13 May	2 May 20 May	8 Jul	17 Jan 18 Jul	6 Aug	21 Jul 16 Aug
Oats	2,980	19.4	11.2	10 Apr 20 May				11.2 15 Aug 5 Sep
Winter Wheat	776	40.4	11.2	25 Apr 15 Jun				
Total 1936 Area Acres	17,012	9.1						

(Continued)

Appendix B. Averages in 1956 and Available Phenological Dates (Approximate) for Selected Crops in the Principal Agricultural Regions of the Soviet Union (continued)

Crop	1956 Average (1,000 ha.)	Percent of Total	Rate of Growth		Rate of Maturity		Rate of Harvest	
			Average	Minimum	Average	Minimum	Average	Minimum
<u>Central Asian Soil Zone</u>								
Winter Rye	2,015	16.4			27 May	14 May 5 June	23 Jul	12 Jul 17 Aug
Winter Wheat	978 2/	5.3			15 June	2 June 28 June	24 Jul	14 Jul 1 Aug
Spring Wheat	2,120	4.3			20 June	15 June 27 June	31 Jul	18 Jul 22 Aug
Spring Barley	806	7.2	24 Apr	12 Apr	19 June	27 May 8 June	25 Jul	7 Jul 15 Aug
Maize	1,433	0.7	13 Apr	27 Mar 25 May			22 Aug	15 Aug 5 Sep
Sugar Beets	405	20.2	22	10 Apr 20 May				
Sunflowers	636	15.2	22	1 Apr 15 May				
Total 1956 Sown Area	15,795	0.1	22	25 Mar 1 June				

(Continued)

Appendix B. Acreage in 1956 and Available Phenological Dates (Approximate) for Selected Crops in the Principal Agricultural Regions of the Soviet Union 2/ (continued)

Crop	1956 Acreage (1,000 ha.)	Percent of USSR Total	Date Down		Date Hatched		Date Ripe			
			Average	Extremes	Average	Extremes	Average	Extremes		
Barley										
Winter Rye	2,060	11.2			22 May	1 May	3 Jun	10 Jul	1 Aug	
Winter Wheat	8,445 b/	46.3			5 Jun	17 May	26 Jun	13 Jul	21 Aug	
Spring Barley	4,210	21.6			13 Jun	27 May	28 Jun	17 Jul	1 Aug	
Oats	1,536	10.3	18 Apr	13 May	23 May					
Flour Rye	213	11.1	22	15 May	1 May			22	10 Jul	5 Aug
Major Rye	1,873	63.4	22	25 Apr	15 Jun					
Buckwheat	1,413	26.7	22	1 Apr	25 May					
Total 1956 Crop Area	22,586	16.7	22	25 May	1 Jun					

(Continued)

Appendix B. Average in 1956 and Available Phenological Dates (Approximate) for Selected Crops in the Principal Agricultural Regions of the Soviet Union ^{2/} (continued)

Crop	1956 Average		Date Sown		Date Harvested		Date Ripe	
	(1,000 ha.)	Percent of Total	Average	Extremes	Average	Extremes	Average	Extremes
North Caucasus								
Winter wheat	4,256 ^{2/}	27.1			5 Jun	19 May	18 Jun	19 Jun 6 Aug
Spring wheat	1,087	2.1	10 Apr	28 Mar	14 Jun	5 Jun	20 Jun	7 Jul 2 Aug
Spring barley	1,493	13.3	16 Apr	25 Mar	22 May	22 May	23 Jun	12 Jul 10 Aug
Oats	255	1.7	22 Apr	20 Mar	22 May	22 May	22 Jun	10 Jul 23 Jul
Barley	1,076	23.8	22 Apr	25 Mar	22 May	22 May	22 Jun	10 Jul 23 Jul
Total 1956 Sown Area	15,659	8.1						

(Continued)

Appendix B. Acreage in 1956 and Available Phenological Dates (Approximate) for Selected Crops in the Principal Agricultural Regions of the Soviet Union/ (continued)

Crop	1956 Acreage (1,000 ha.)	Percent of USSR Total	Date Began		Date Harvested		Date Began	
			Average	Extremes	Average	Extremes	Average	Extremes
Wheat								
Winter	2,344	12.7			1 Jan	19 May	11 Jun	15 Jul
Winter	360 2/	2.0			8 Jan	28 May	21 Jun	11 Jul
Spring	5,646	11.5			24 Jan	30 May	15 Jul	1 Aug
Spring	1,046	5.3			22 Jan	19 Jun	26 Jun	22 Jul
Oats	643	4.3						2 Aug
Barley	540	10.2						5 Aug
Total 1956 Sown Area	16,202	8.3						

(Continued)

AVERAGE DATES AND EXTREMES OF SOWING, HEADING, AND RIPENING OF SPRING BARLEY (LOCAL VARIETIES) IN THE SOVIET UNION

LEGEND

- Sown: (Symbol: a small circle with a dot)
- Headed: (Symbol: a small circle with a cross)
- Ripe: (Symbol: a small circle with a triangle)
- Dates outside parentheses represent average date.
- Dates in parentheses represent extremes.

Region	City	Sown	Headed	Ripe
Northern Region	Khibiny	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
	Karelia	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
Central Region	Leningrad	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
	Moscow	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
Southern Region	Ukraine	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
	Belarus	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
Eastern Region	Far East	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
	Amur	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
Western Region	Poland	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)
	Czechoslovakia	May 21 (18-29)	June 10 (7-21)	June 21 (18-29)

LEGEND	
Sown	-----
Headed	-----
Ripe	-----
Dates outside parentheses represent average date.	
Dates in parentheses represent extremes.	

1. <u>NAME</u> 2. <u>ADDRESS</u> 3. <u>CITY</u> 4. <u>STATE</u> 5. <u>ZIP</u> 6. <u>PHONE</u> 7. <u>TELETYPE</u> 8. <u>TELEFAX</u> 9. <u>TELEMAIL</u> 10. <u>TELEFAX</u> 11. <u>TELEMAIL</u> 12. <u>TELEFAX</u> 13. <u>TELEMAIL</u> 14. <u>TELEFAX</u> 15. <u>TELEMAIL</u> 16. <u>TELEFAX</u> 17. <u>TELEMAIL</u> 18. <u>TELEFAX</u> 19. <u>TELEMAIL</u> 20. <u>TELEFAX</u> 21. <u>TELEMAIL</u> 22. <u>TELEFAX</u> 23. <u>TELEMAIL</u> 24. <u>TELEFAX</u> 25. <u>TELEMAIL</u> 26. <u>TELEFAX</u> 27. <u>TELEMAIL</u> 28. <u>TELEFAX</u> 29. <u>TELEMAIL</u> 30. <u>TELEFAX</u> 31. <u>TELEMAIL</u> 32. <u>TELEFAX</u> 33. <u>TELEMAIL</u> 34. <u>TELEFAX</u> 35. <u>TELEMAIL</u> 36. <u>TELEFAX</u> 37. <u>TELEMAIL</u> 38. <u>TELEFAX</u> 39. <u>TELEMAIL</u> 40. <u>TELEFAX</u> 41. <u>TELEMAIL</u> 42. <u>TELEFAX</u> 43. <u>TELEMAIL</u> 44. <u>TELEFAX</u> 45. <u>TELEMAIL</u> 46. <u>TELEFAX</u> 47. <u>TELEMAIL</u> 48. <u>TELEFAX</u> 49. <u>TELEMAIL</u> 50. <u>TELEFAX</u> 51. <u>TELEMAIL</u> 52. <u>TELEFAX</u> 53. <u>TELEMAIL</u> 54. <u>TELEFAX</u> 55. <u>TELEMAIL</u> 56. <u>TELEFAX</u> 57. <u>TELEMAIL</u> 58. <u>TELEFAX</u> 59. <u>TELEMAIL</u> 60. <u>TELEFAX</u> 61. <u>TELEMAIL</u> 62. <u>TELEFAX</u> 63. <u>TELEMAIL</u> 64. <u>TELEFAX</u> 65. <u>TELEMAIL</u> 66. <u>TELEFAX</u> 67. <u>TELEMAIL</u> 68. <u>TELEFAX</u> 69. <u>TELEMAIL</u> 70. <u>TELEFAX</u> 71. <u>TELEMAIL</u> 72. <u>TELEFAX</u> 73. <u>TELEMAIL</u> 74. <u>TELEFAX</u> 75. <u>TELEMAIL</u> 76. <u>TELEFAX</u> 77. <u>TELEMAIL</u> 78. <u>TELEFAX</u> 79. <u>TELEMAIL</u> 80. <u>TELEFAX</u> 81. <u>TELEMAIL</u> 82. <u>TELEFAX</u> 83. <u>TELEMAIL</u> 84. <u>TELEFAX</u> 85. <u>TELEMAIL</u> 86. <u>TELEFAX</u> 87. <u>TELEMAIL</u> 88. <u>TELEFAX</u> 89. <u>TELEMAIL</u> 90. <u>TELEFAX</u> 91. <u>TELEMAIL</u> 92. <u>TELEFAX</u> 93. <u>TELEMAIL</u> 94. <u>TELEFAX</u> 95. <u>TELEMAIL</u> 96. <u>TELEFAX</u> 97. <u>TELEMAIL</u> 98. <u>TELEFAX</u> 99. <u>TELEMAIL</u> 100. <u>TELEFAX</u> 101. <u>TELEMAIL</u> 102. <u>TELEFAX</u> 103. <u>TELEMAIL</u> 104. <u>TELEFAX</u> 105. <u>TELEMAIL</u> 106. <u>TELEFAX</u> 107. <u>TELEMAIL</u> 108. <u>TELEFAX</u> 109. <u>TELEMAIL</u> 110. <u>TELEFAX</u> 111. <u>TELEMAIL</u> 112. <u>TELEFAX</u> 113. <u>TELEMAIL</u> 114. <u>TELEFAX</u> 115. <u>TELEMAIL</u> 116. <u>TELEFAX</u> 117. <u>TELEMAIL</u> 118. <u>TELEFAX</u> 119. <u>TELEMAIL</u> 120. <u>TELEFAX</u> 121. <u>TELEMAIL</u> 122. <u>TELEFAX</u> 123. <u>TELEMAIL</u> 124. <u>TELEFAX</u> 125. <u>TELEMAIL</u> 126. <u>TELEFAX</u> 127. <u>TELEMAIL</u> 128. <u>TELEFAX</u> 129. <u>TELEMAIL</u> 130. <u>TELEFAX</u> 131. <u>TELEMAIL</u> 132. <u>TELEFAX</u> 133. <u>TELEMAIL</u> 134. <u>TELEFAX</u> 135. <u>TELEMAIL</u> 136. <u>TELEFAX</u> 137. <u>TELEMAIL</u> 138. <u>TELEFAX</u> 139. <u>TELEMAIL</u> 140. <u>TELEFAX</u> 141. <u>TELEMAIL</u> 142. <u>TELEFAX</u> 143. <u>TELEMAIL</u> 144. <u>TELEFAX</u> 145. <u>TELEMAIL</u> 146. <u>TELEFAX</u> 147. <u>TELEMAIL</u> 148. <u>TELEFAX</u> 149. <u>TELEMAIL</u> 150. <u>TELEFAX</u> 151. <u>TELEMAIL</u> 152. <u>TELEFAX</u> 153. <u>TELEMAIL</u> 154. <u>TELEFAX</u> 155. <u>TELEMAIL</u> 156. <u>TELEFAX</u> 157. <u>TELEMAIL</u> 158. <u>TELEFAX</u> 159. <u>TELEMAIL</u> 160. <u>TELEFAX</u> 161. <u>TELEMAIL</u> 162. <u>TELEFAX</u> 163. <u>TELEMAIL</u> 164. <u>TELEFAX</u> 165. <u>TELEMAIL</u> 166. <u>TELEFAX</u> 167. <u>TELEMAIL</u> 168. <u>TELEFAX</u> 169. <u>TELEMAIL</u> 170. <u>TELEFAX</u> 171. <u>TELEMAIL</u> 172. <u>TELEFAX</u> 173. <u>TELEMAIL</u> 174. <u>TELEFAX</u> 175. <u>TELEMAIL</u> 176. <u>TELEFAX</u> 177. <u>TELEMAIL</u> 178. <u>TELEFAX</u> 179. <u>TELEMAIL</u> 180. <u>TELEFAX</u> 181. <u>TELEMAIL</u> 182. <u>TELEFAX</u> 183. <u>TELEMAIL</u> 184. <u>TELEFAX</u> 185. <u>TELEMAIL</u> 186. <u>TELEFAX</u> 187. <u>TELEMAIL</u> 188. <u>TELEFAX</u> 189. <u>TELEMAIL</u> 190. <u>TELEFAX</u> 191. <u>TELEMAIL</u> 192. <u>TELEFAX</u> 193. <u>TELEMAIL</u> 194. <u>TELEFAX</u> 195. <u>TELEMAIL</u> 196. <u>TELEFAX</u> 197. <u>TELEMAIL</u> 198. <u>TELEFAX</u> 199. <u>TELEMAIL</u> 200. <u>TELEFAX</u> 201. <u>TELEMAIL</u> 202. <u>TELEFAX</u> 203. <u>TELEMAIL</u> 204. <u>TELEFAX</u> 205. <u>TELEMAIL</u> 206. <u>TELEFAX</u> 207. <u>TELEMAIL</u> 208. <u>TELEFAX</u> 209. <u>TELEMAIL</u> 210. <u>TELEFAX</u> 211. <u>TELEMAIL</u> 212. <u>TELEFAX</u> 213. <u>TELEMAIL</u> 214. <u>TELEFAX</u> 215. <u>TELEMAIL</u> 216. <u>TELEFAX</u> 217. <u>TELEMAIL</u> 218. <u>TELEFAX</u> 219. <u>TELEMAIL</u> 220. <u>TELEFAX</u> 221. <u>TELEMAIL</u> 222. <u>TELEFAX</u> 223. <u>TELEMAIL</u> 224. <u>TELEFAX</u> 225. <u>TELEMAIL</u> 226. <u>TELEFAX</u> 227. <u>TELEMAIL</u> 228.
--

AVERAGE DATES AND EXTREMES OF SOWING, HEADING, AND RIPENING OF SPRING BARLEY (LOCAL VARIETIES) AT AGRICULTURAL EXPERIMENT STATIONS IN THE SOVIET UNION

(Arranged from earliest to latest average ripening dates)

LOCATION	LAT. (N)	LONG. (E)	DATE SOWN			DATE HEADED			DATE RIPE		
			AVERAGE	EXTREMES		AVERAGE	EXTREMES		AVERAGE	EXTREMES	
Shkhabad, Turkmen S.S.R.	37°57'	58°23'	Apr. 1	(11)	Mar. 10-Apr. 12	May 29	(11)	May 26-June 16	June 20	(11)	June 21-July 17
Tilok, Ukrainian S.S.R.	46°07'	22°51'	Mar. 30		Mar. 13-May 3	June 7		June 2-June 15	July 2		July 4-July 9
Idessa, Ukrainian S.S.R.	46°29'	30°44'	Apr. 6		Mar. 21-Apr. 15	June 8		June 1-June 15	July 7		July 1-July 13
erson, Ukrainian S.S.R.	46°38'	32°37'	Apr. 2		Mar. 23-Apr. 10	June 15		June 10-June 20	...do...		...do...
Lejandrovsk, Ukrainian S.S.R.	47°49'	35°11'	Apr. 1		Mar. 24-Apr. 22	June 4		May 25-June 16	July 9		July 4-July 14
Lizhneye Donantse, Ukrainian S.S.R.	48°38'	22°20'	Apr. 8		Mar. 26-Apr. 8	June 2		May 22-June 12	July 12		July 2-July 22
Velikaya Bakta, Ukrainian S.S.R.	48°10'	22°40'	Apr. 3		Mar. 26-Apr. 8	June 2		May 22-June 12	July 12		July 9-July 18
Velikiye Komary, Ukrainian S.S.R.	48°14'	22°58'	Apr. 10		Apr. 4-Apr. 13	June 13		May 27-June 25	July 13		July 12-July 30
Rostov-on-the-Don, R.S.F.S.R.	47°13'	39°43'	Apr. 19		Apr. 29-Apr. 26	(11)		June 5-June 22	July 20		July 13-Aug. 1
Ploty, Moldavian S.S.R.	47°57'	29°10'	Apr. 2		Mar. 24-Apr. 27	June 18		June 19-June 26	July 21		July 21-July 25
Saratov, R.S.F.S.R.	51°32'	46°01'	Apr. 23		Apr. 3-May 8	June 22		June 18-June 28	July 22		July 13-Aug. 1
Velikiy Bychkov, Ukrainian S.S.R.	47°59'	24°02'	Apr. 17		Apr. 10-Apr. 29	June 21		May 27-July 8	July 23		July 13-Aug. 1
Poltava, Ukrainian S.S.R.	49°35'	34°34'	Apr. 23		Mar. 27-May 23	June 16		May 27-July 8	...do...		July 7-Aug. 6
Stavropol: West Field, R.S.F.S.R.	45°02'	41°54'	Apr. 11		Mar. 28-May 3	n.a.		June 6-July 11	July 25		July 17-Aug. 3
Stavropol: East Field, R.S.F.S.R.	45°02'	41°54'	Apr. 24		Mar. 25-May 18	June 25		June 6-July 11	July 26		July 13-Aug. 10
Kharkov, Ukrainian S.S.R.	49°58'	36°09'	Apr. 13		Apr. 2-Apr. 22	n.a.		June 22-June 28	July 31		July 18-Aug. 15
Zyryanovsk, Kazakh S.S.R.	49°44'	84°18'	May 20		May 16-May 22	(11)		June 22-June 28	Aug. 3		July 21-Aug. 12
Tursk, White Russian S.S.R.	53°02'	30°13'	May 7		Apr. 26-May 15	n.a.		June 17-July 3	...do...		July 24-Aug. 12
Barnaul, R.S.F.S.R.	53°20'	83°48'	May 21		May 10-May 26	(11)		June 17-July 3	...do...		July 28-Aug. 11
Yakutsk, R.S.F.S.R.	61°29'	129°09'	May 20		May 16-May 24	July 6		July 2-July 9	Aug. 4		July 27-Aug. 12
Nizhniye Vorota, Ukrainian S.S.R.	48°46'	23°06'	Apr. 16		Mar. 31-May 2	June 24		June 20-June 29	Aug. 6		July 22-Aug. 13
Krivoshchekovo, R.S.F.S.R.	54°59'	82°52'	May 17		May 14-May 23	(11)		June 22-July 7	...do...		July 25-Aug. 15
Moscow, R.S.F.S.R.	55°50'	37°33'	May 10		May 5-May 20	July 1		June 27-July 5	Aug. 7		Aug. 1-Aug. 13
Leninsk-Kuznetskiy, R.S.F.S.R.	54°43'	86°13'	May 17		May 12-May 23	(11)		June 13-July 3	...do...		July 29-Aug. 18
Kirov, R.S.F.S.R.	58°36'	49°41'	May 13		May 2-May 23	(11)		June 13-July 3	Aug. 8		July 28-Aug. 23
Bel-Agach, Kazakh S.S.R.	50°48'	80°44'	May 25		May 16-June 1	(11)		June 13-July 3	...do...		July 24-Aug. 20
Borovye Ozera R.S.F.S.R.	51°45'	80°20'	May 24		May 17-May 30	(11)		July 3-July 19	Aug. 9		Aug. 6-Aug. 13
Urkul, R.S.F.S.R.	52°41'	84°37'	May 16		May 12-May 20	(11)		July 3-July 19	Aug. 10		July 31-Aug. 22
Stalinsk, R.S.F.S.R.	53°46'	87°11'	May 22		May 18-May 29	(11)		July 5-July 20	Aug. 11		Aug. 1-Aug. 20
Krasnoyarsk, R.S.F.S.R.	56°08'	93°00'	May 16		May 13-May 19	July 13		July 13-July 18	Aug. 15		Aug. 2-Aug. 22
Zapolye, R.S.F.S.R.	59°23'	29°30'	May 26		May 21-May 29	July 16		July 13-July 18	Aug. 20		Aug. 3-Sept. 26
Leningrad, R.S.F.S.R.	59°56'	30°16'	May 27		May 9-June 17	n.a.		July 13-July 18	Aug. 25		Aug. 8-Sept. 14
Khibiny, R.S.F.S.R.	67°44'	33°02'	June 4		May 27	(11)		July 13-July 18	Sept. 1		Aug. 10-Sept. 9
Minsk, White Russian S.S.R.	53°54'	27°33'	May 10		Apr. 25-May 23	n.a.		July 13-July 18	Sept. 1		Aug. 10-Sept. 9

Source: Based on data as shown in Tables 67 through 90 and Table 93. *Indicates stations for which day-degree summations were computed for individual years. See Tables 67 through 90.

†Not computed because of small number of cases.
n.a. - not available.

Figure 4




TABLE 2
AVERAGE DATES AND EXTREMES OF SOWING, HEADING, AND RIPENING OF WINTER RYE (LOCAL VARIETIES)
AT AGRICULTURAL EXPERIMENT STATIONS IN THE SOVIET UNION

(Arranged from earliest to latest average ripening dates)

LOCATION	LAT. (N)	LONG. (E)	DATE SOWN		DATE HEADED		DATE RIFE	
			AVERAGE	EXTREMES	AVERAGE	EXTREMES	AVERAGE	EXTREMES
*Kherson, Ukrainian S.S.R.	46°38'	32°37'	Sept. 22	Aug. 22 - Oct. 29	May 11	May 3 - May 24	June 26	June 16 - July 3
*Vilok, Ukrainian S.S.R.	48°07'	22°51'	Sept. 17	Sept. 1 - Oct. 21	May 20	May 13 - May 28	June 28	June 18 - July 14
*Ploty, Moldavian S.S.R.	47°57'	29°10'	Sept. 3	Aug. 25 - Sept. 25	May 13	May 6 - May 24	July 3	June 25 - July 13
*Aleksandropol, Ukrainian S.S.R.	48°01'	34°27'	Sept. 5	Aug. 22 - Sept. 23	May 18	May 11 - May 27	July 5	June 25 - July 17
*Poltava, Ukrainian S.S.R.	49°35'	34°34'	Aug. 30	Aug. 22 - Sept. 22	May 13	May 1 - May 24	July 7	July 1 - July 13
*Nizhneye Domanintse, Ukrainian S.S.R.	48°38'	22°20'	Oct. 5	Sept. 18 - Oct. 20	May 20	May 12 - May 28	July 7	July 1 - July 11
*Nalchik, R.S.F.S.R.	43°27'	43°35'	Sept. 18	Sept. 15 - Sept. 21	May 14	n.a.	July 5	July 7 - July 10
*Migeyev, Ukrainian S.S.R.	48°02'	30°57'	(1)	(1)	May 16	May 10 - May 21	July 10	July 5 - July 14
*Krasnyy Kut, R.S.F.S.R.	50°58'	46°30'	Sept. 15	Aug. 25 - Oct. 23	May 29	May 20 - June 3	July 11	July 3 - July 16
*Saratov, R.S.F.S.R.	51°32'	46°01'	Aug. 24	Aug. 21 - Aug. 27	May 27	May 19 - June 7	July 11	July 2 - July 15
*Kazan, R.S.F.S.R.	55°47'	49°08'	Aug. 15	Aug. 10 - Aug. 18	May 31	May 26 - June 7	July 11	July 8 - July 15
*Kharkov, Ukrainian S.S.R.	49°58'	36°09'	Aug. 28	Aug. 20 - Oct. 15	n.a.	---	July 12	June 28 - July 22
*Bezenchuk, R.S.F.S.R.	53°00'	49°28'	Aug. 26	Aug. 20 - Sept. 5	June 1	May 23 - June 11	July 12	July 8 - July 17
*Velikiy Bychkov, Ukrainian S.S.R.	47°59'	24°02'	Oct. 25	Oct. 20 - Oct. 26	May 27	May 25 - June 1	July 12	July 10 - July 31
*Bogoroditskoye, R.S.F.S.R.	51°10'	37°21'	n.a.	---	(1)	May 14 - May 26	July 13	July 13 - July 24
*Kon-Kolodez, R.S.F.S.R.	52°08'	39°10'	Aug. 24	Aug. 22 - Aug. 26	May 26	May 21 - June 1	July 13	July 15 - July 22
*Uyutnoye, R.S.F.S.R.	52°04'	35°05'	Sept. 1	Aug. 29 - Sept. 2	---	May 17 - June 5	July 20	July 11 - July 30
*Belebey, R.S.F.S.R.	53°55'	54°38'	Aug. 14	(1)	June 1	May 28 - June 6	---	(1)
*Novo-Urensk, R.S.F.S.R.	54°25'	48°11'	Aug. 27	Aug. 17 - Sept. 11	June 6	June 5 - June 8	---	July 13 - July 27
*Ufa, R.S.F.S.R.	54°43'	55°56'	Aug. 20	Aug. 7 - Sept. 5	June 8	May 30 - June 26	---	July 5 - Aug. 9
*Dovzhik, Ukrainian S.S.R.	51°37'	31°01'	Aug. 30	Aug. 27 - Aug. 31	May 31	May 27 - June 4	July 21	July 10 - Aug. 1
*Bukhovoye, R.S.F.S.R.	53°03'	39°58'	Aug. 16	Aug. 13 - Aug. 21	May 27	May 18 - June 2	---	July 14 - Aug. 17
*Itmanovo, R.S.F.S.R.	55°08'	44°45'	Aug. 24	Aug. 15 - Sept. 4	May 29	May 22 - June 8	---	July 10 - Aug. 5
*Zyranovsk, Kazakh S.S.R.	49°44'	84°18'	Aug. 12	Aug. 8 - Aug. 18	(1)	(1)	July 22	July 14 - Aug. 2
*Pogozheye, R.S.F.S.R.	51°36'	37°16'	(1)	(1)	June 8	June 2 - June 14	---	July 18 - July 27
*Marina Gorka, White Russian S.S.R.	53°26'	28°18'	Sept. 6	Sept. 4 - Sept. 7	May 26	May 17 - June 4	---	July 13 - July 31
*Bogoroditsk, R.S.F.S.R.	53°47'	38°07'	Aug. 20	Aug. 11 - Sept. 3	May 31	May 20 - June 11	---	July 5 - Aug. 11
*Minsk, White Russian S.S.R.	53°54'	27°33'	n.a.	---	May 25	May 14 - May 31	---	July 17 - July 30
*Serpukhov, R.S.F.S.R.	54°54'	37°24'	Aug. 20	n.a.	May 30	n.a.	---	n.a.
*Shatilov, R.S.F.S.R.	53°00'	37°23'	---	Aug. 12 - Aug. 29	---	May 19 - June 7	July 23	July 8 - Aug. 1
*Belyy, R.S.F.S.R.	55°50'	32°55'	n.a.	---	May 28	May 18 - June 8	---	July 10 - Aug. 4
*Chokino, R.S.F.S.R.	52°14'	41°38'	Aug. 18	Aug. 7 - Aug. 27	(1)	May 31 - June 4	July 24	July 17 - July 28
*Moscow, R.S.F.S.R.	55°50'	37°33'	Sept. 2	Aug. 27 - Sept. 7	May 31	May 11 - June 15	---	June 29 - Aug. 12
*Kirov, R.S.F.S.R.	58°36'	49°41'	Aug. 22	Aug. 20 - Aug. 24	May 30	May 26 - June 4	---	July 23 - July 25
*Bobruysk, White Russian S.S.R.	53°08'	29°15'	Sept. 14	Sept. 10 - Sept. 18	May 28	May 17 - June 9	July 25	July 12 - Aug. 7
*Kalinin, R.S.F.S.R.	56°50'	35°55'	n.a.	---	n.a.	---	---	July 15 - Aug. 3
*Voronezh, R.S.F.S.R.	51°36'	38°59'	Aug. 19	Aug. 17 - Aug. 21	May 29	May 26 - May 31	July 27	July 24 - July 29
*Borovye Ozera, R.S.F.S.R.	51°45'	80°20'	Aug. 15	Aug. 5 - Aug. 21	n.a.	---	---	July 23 - Aug. 1
*Pavlovskiy Posad, R.S.F.S.R.	55°47'	38°40'	Aug. 20	n.a.	May 30	n.a.	---	n.a.
*Pskov, R.S.F.S.R.	57°33'	28°19'	n.a.	---	(1)	May 29 - June 9	---	July 20 - Aug. 8
*Boraul, R.S.F.S.R.	53°20'	83°48'	Aug. 21	Aug. 12 - Sept. 3	(1)	June 11 - June 28	July 28	July 21 - Aug. 2
*Gibitsy, T.S.F.S.R.	54°58'	41°05'	---	Aug. 10 - Aug. 29	May 28	May 17 - June 3	---	July 14 - Aug. 14
*Istra, R.S.F.S.R.	55°55'	36°55'	Aug. 22	Aug. 15 - Sept. 3	May 30	May 16 - June 10	---	July 15 - Aug. 15
*Utkul, R.S.F.S.R.	52°41'	84°37'	Aug. 17	Aug. 13 - Aug. 21	(1)	(1)	July 14	July 24 - Aug. 1
*Tursk, White Russian S.S.R.	53°02'	30°13'	Sept. 11	Aug. 27 - Oct. 6	June 10	June 3 - June 21	---	July 16 - Aug. 6
*Stalinsk, R.S.F.S.R.	53°46'	87°11'	Aug. 18	Aug. 12 - Aug. 27	(1)	(1)	---	July 20 - Aug. 5
*Kungur, R.S.F.S.R.	57°25'	56°56'	Aug. 24	Aug. 9 - Aug. 31	June 7	June 3 - June 10	---	July 22 - Aug. 4
*Nerekhta, R.S.F.S.R.	57°28'	40°35'	n.a.	---	May 29	May 20 - June 4	---	July 15 - Aug. 15
*Leninsk-Kuznetskiy, R.S.F.S.R.	54°43'	86°13'	Aug. 19	Aug. 17 - Aug. 25	(1)	June 6 - June 14	July 28	July 22 - Aug. 9
*Krivoshchevovo, R.S.F.S.R.	54°59'	82°52'	Aug. 16	Aug. 13 - Aug. 25	(1)	(1)	---	---
*Batishchevo, R.S.F.S.R.	55°12'	33°23'	n.a.	---	May 29	May 22 - June 4	---	July 20 - Aug. 15
*Yuryev-Polskiy, R.S.F.S.R.	56°30'	39°40'	n.a.	---	June 16	June 9 - June 25	---	July 18 - Aug. 17
*Koblukovo, R.S.F.S.R.	56°50'	36°12'	n.a.	---	May 31	May 17 - June 10	July 30	July 25 - Aug. 16
*Soligalich, R.S.F.S.R.	59°05'	42°17'	n.a.	---	n.a.	---	---	July 20 - Aug. 9
*Putilovo, R.S.F.S.R.	59°52'	31°25'	n.a.	---	June 6	May 26 - June 19	---	July 19 - Aug. 14
*Bel-Agach, Kazakh S.S.R.	50°48'	60°44'	Aug. 21	Aug. 14 - Aug. 30	n.a.	---	Aug. 1	July 26 - Aug. 9
*Klin, R.S.F.S.R.	56°21'	36°45'	Aug. 20	n.a.	June 5	n.a.	Aug. 2	n.a.
*Moletov, R.S.F.S.R.	58°01'	56°16'	Aug. 16	Aug. 10 - Aug. 22	June 8	June 5 - June 11	---	July 31 - Aug. 4
*Shimsk, R.S.F.S.R.	58°12'	30°40'	Sept. 1	Aug. 19 - Sept. 15	(1)	(1)	---	July 20 - Aug. 8
*Nikolsk, R.S.F.S.R.	59°22'	45°27'	n.a.	---	June 5	June 2 - June 6	Aug. 6	July 29 - Aug. 14
*Bazlovo, R.S.F.S.R.	56°30'	30°01'	Sept. 6	Sept. 1 - Sept. 11	June 3	May 21 - June 15	Aug. 7	July 26 - Aug. 19
*Cherepovets, R.S.F.S.R.	59°07'	37°56'	n.a.	---	June 6	June 2 - June 9	---	July 30 - Aug. 14
*Leningrad, R.S.F.S.R.	59°56'	30°16'	Aug. 26	Aug. 22 - Aug. 31	June 7	June 3 - June 10	Aug. 8	Aug. 3 - Aug. 17
*Cherdyn, R.S.F.S.R.	60°24'	56°21'	---	Aug. 12 - Aug. 18	June 20	June 10 - June 27	Aug. 14	Aug. 8 - Aug. 21

Source: Based on data as shown in Tables 22 through 52 and Table 53, and on data from agricultural experiment stations in the Soviet Union between 1914 and 1940. Indicates stations for which a 6-year summary is available; data for individual years. See Tables 22 through 52.

*Not computed because of small number of cases.
n.a. = not available.

AVERAGE DATES AND EXTREMES OF SOWING, HEADING, AND RIPENING OF WINTER WHEAT IN THE U.S.S.R.

LEGEND

- Sown - - - - -
- Headed - - - - -
- Ripe - - - - -
- Dates outside parenthesis represent average dates
- Dates in parenthesis represent extremes

The map displays the following data points for various locations in the U.S.S.R. (Coordinates are approximate):

- Moscow:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Leningrad:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Kazakhstan:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Ukraine:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Belarus:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Poland:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Czech Republic:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Slovakia:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Hungary:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Romania:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Bulgaria:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Greece:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Turkey:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Iran:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- India:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- China:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Japan:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- USA:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Canada:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- South America:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Africa:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Europe:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Asia:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Oceania:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)
- Antarctica:** Sown (Aug 15/25-30/35), Headed (Jul 15/25-30/35), Ripe (Jun 15/25-30/35)

LEGEND

Same -----
Headed -----
Ripe -----
Dates outside parenthesis
represent average dates.
Dates in parenthesis represent
extremes.

TABLE 2

AVERAGE DATES AND EXTREMES OF SOWING, HEADING, AND RIPENING OF WINTER WHEAT¹ AT AGRICULTURAL EXPERIMENT STATIONS IN THE SOVIET UNION

(Arranged from earliest to latest average ripening dates)

LOCATION	LAT. (N)	LONG. (E)	DATE SOWN			DATE HEADED			DATE RIPE		
			AVERAGE	EXTREMES	AVERAGE	AVERAGE	EXTREMES	AVERAGE	AVERAGE	EXTREMES	EXTREMES
Asikhabad, Turkmen S.S.R.	37° 57'	58° 23'	n.a.	—	—	—	—	—	June 6	June 5—June	June 5—June
Kamashi, Uzbek S.S.R.	38° 51'	65° 23'	Nov. 3	Oct. 10—Nov. 27	n.a.	May 8	Apr. 26—May 20	June 10	June 10	May 31—June	May 31—June
Krasnovodopadsk, Uzbek S.S.R.	40° 57'	69° 31'	Nov. 13	Nov. 9—Nov. 17	May 18	May 18	May 8—May 28	June 12	June 12	June 8—June	June 8—June
Stalinabad, Tadzhik S.S.R.	38° 30'	68° 45'	Nov. 28	Nov. 11—Dec. 22	May 22	May 22	May 13—June 1	June 15	June 15	June 8—June	June 8—June
Katta-Kurgan, Uzbek S.S.R.	39° 55'	66° 15'	Nov. 13	Nov. 4—Nov. 22	May 14	May 14	Apr. 30—May 28	June 16	June 16	June 5—June	June 5—June
Milyutin, Uzbek S.S.R.	40° 02'	67° 36'	Nov. 5	Oct. 28—Nov. 13	May 25	May 25	May 16—June 3	June 22	June 22	June 16—June	June 16—June
Khasavyurt, R.S.F.S.R.	43° 24'	46° 10'	Oct. 12	Oct. 5—Oct. 19	June 1	June 1	May 24—June 10	June 26	June 26	June 18—July	June 18—July
Kuban, R.S.F.S.R.	45° 00'	37° 30'	Oct. 5	Sept. 27—Oct. 14	May 30	May 30	May 21—June 7	June 29	June 29	June 22—July	June 22—July
*Kherson, Ukrainian S.S.R.	46° 38'	32° 37'	Sept. 5	Aug. 23—Sept. 28	May 27	May 27	May 19—June 2	...do...	...do...	June 21—July	June 21—July
Nalchik, R.S.F.S.R.	43° 34'	44° 04'	Oct. 1	Sept. 17—Oct. 14	June 2	June 2	May 25—June 10	June 30	June 30	June 22—July	June 22—July
*Krasnodar, R.S.F.S.R.	45° 02'	38° 56'	Oct. 21	Oct. 2—Nov. 3	June 1	June 1	May 19—June 11	...do...	...do...	June 19—July	June 19—July
Budennovsk, R.S.F.S.R.	44° 46'	44° 12'	Oct. 2	Sept. 27—Oct. 6	May 30	May 30	May 21—June 7	...do...	...do...	June 21—July	June 21—July
Frunze, Kirgiz S.S.R.	42° 54'	74° 36'	Sept. 29	Sept. 25—Oct. 3	May 26	May 26	May 23—May 30	July 1	July 1	June 26—July	June 26—July
Krasnogvardeyskoye, R.S.F.S.R.	45° 30'	34° 18'	Oct. 13	Sept. 24—Nov. 3	...do...	...do...	May 15—June 7	...do...	...do...	June 17—July	June 17—July
*Odessa, Ukrainian S.S.R.	46° 29'	30° 44'	Sept. 19	Aug. 20—Oct. 15	May 25	May 25	May 17—June 3	July 4	July 4	June 24—July	June 24—July
*Rostov-Na-Donu, R.S.F.S.R.	47° 13'	39° 43'	Sept. 15	Sept. 9—Sept. 27	May 31	May 31	May 24—June 16	...do...	...do...	June 27—July	June 27—July
*Krasnyy Kut, R.S.F.S.R.	50° 58'	46° 30'	Aug. 31	Aug. 28—Sept. 5	June 9	June 9	June 3—June 16	July 7	July 7	June 29—July	June 29—July
Alma-Ata, Kazakh S.S.R.	43° 15'	76° 55'	Sept. 19	Sept. 1—Oct. 10	June 6	June 6	May 31—June 8	July 8	July 8	July 5—July	July 5—July
*Sinelnikovo, Ukrainian S.S.R.	48° 21'	35° 26'	Sept. 7	Sept. 3—Sept. 11	...do...	...do...	May 29—June 17	...do...	...do...	June 30—July	June 30—July
Stalingrad, R.S.F.S.R.	48° 42'	44° 31'	Sept. 10	Sept. 7—Sept. 14	...do...	...do...	May 29—June 14	...do...	...do...	July 1—July	July 1—July
Tselinna, R.S.F.S.R.	46° 25'	41° 00'	Sept. 24	Sept. 17—Oct. 1	June 3	June 3	May 27—June 9	July 10	July 10	July 2—July	July 2—July
*Zherebkovo, Ukrainian S.S.R.	47° 51'	29° 53'	Sept. 10	Sept. 2—Sept. 14	June 7	June 7	June 2—June 13	...do...	...do...	July 3—July	July 3—July
*Kamyshin, R.S.F.S.R.	50° 02'	45° 07'	Aug. 30	Aug. 27—Sept. 2	June 10	June 10	June 5—June 15	...do...	...do...	June 30—July	June 30—July
*Kharkov, Ukrainian S.S.R.	49° 58'	36° 09'	Sept. 1	Aug. 25—Sept. 7	June 7	June 7	May 28—June 17	July 13	July 13	July 5—July	July 5—July
Artemovsk, Ukrainian S.S.R.	48° 35'	38° 02'	Sept. 18	Sept. 7—Sept. 28	June 14	June 14	June 7—June 21	July 14	July 14	July 6—July	July 6—July
*Mironovka, Ukrainian S.S.R.	49° 40'	31° 00'	Sept. 8	Sept. 2—Sept. 15	June 4	June 4	May 27—June 16	...do...	...do...	July 3—July	July 3—July
Gorskiy, R.S.F.S.R.	43° 15'	44° 50'	Oct. 20	Oct. 11—Oct. 29	June 12	June 12	May 30—June 25	July 15	July 15	July 6—July	July 6—July
*Ploty, Moldavian S.S.R.	47° 57'	29° 10'	Aug. 31	Aug. 25—Sept. 5	June 1	June 1	May 18—June 11	...do...	...do...	July 12—July	July 12—July
Temir, Kazakh S.S.R.	49° 06'	57° 10'	Aug. 23	Aug. 20—Aug. 29	n.a.	n.a.	—	...do...	...do...	July 16—July	July 16—July
Uralsk, Kazakh S.S.R.	51° 12'	51° 21'	Sept. 8	Sept. 1—Sept. 14	June 14	June 14	June 9—June 18	...do...	...do...	July 10—July	July 10—July
*Saratov, R.S.F.S.R.	51° 32'	46° 01'	Aug. 30	Aug. 17—Sept. 11	June 9	June 9	May 28—June 21	...do...	...do...	July 4—July	July 4—July
*Poltava, Ukrainian S.S.R.	49° 35'	34° 34'	Aug. 27	Aug. 14—Sept. 28	June 2	June 2	May 18—June 21	July 16	July 16	June 28—July	June 28—July
Chkalov, R.S.F.S.R.	51° 45'	55° 06'	Sept. 2	Aug. 31—Sept. 14	June 7	June 7	May 28—June 17	...do...	...do...	July 14—July	July 14—July
Shterovka, Ukrainian S.S.R.	48° 03'	38° 55'	Sept. 18	Sept. 14—Sept. 20	June 18	June 18	June 8—June 29	July 17	July 17	July 8—July	July 8—July
*Verkhnyachka, Ukrainian S.S.R.	48° 49'	30° 03'	Sept. 10	Aug. 29—Sept. 24	June 8	June 8	May 30—June 18	July 18	July 18	July 8—July	July 8—July
*Bezenchuk, R.S.F.S.R.	53° 00'	49° 28'	Aug. 27	Aug. 23—Sept. 3	June 14	June 14	June 6—June 20	...do...	...do...	July 8—July	July 8—July
*Sumy, Ukrainian S.S.R.	50° 53'	34° 43'	Sept. 10	Sept. 4—Sept. 16	...do...	...do...	June 2—June 25	July 20	July 20	July 4—July	July 4—July
*Nossovka, Ukrainian S.S.R.	50° 56'	31° 25'	Sept. 1	Aug. 20—Sept. 17	n.a.	n.a.	—	...do...	...do...	July 4—Aug	July 4—Aug
*Kamennaya-Steppe, R.S.F.S.R.	51° 03'	40° 42'	Aug. 26	Aug. 23—Sept. 3	June 16	June 16	June 10—June 24	July 21	July 21	July 15—July	July 15—July

Station	Lat.	Long.	Alt.	Year	Month	Day	Temp.	Wind	Humid.	Clouds	Pressure	State	Remarks
Voronezh, R.S.F.S.R.	51° 36'	38° 59'	150	1952	Aug.	26	15.0	1.5	75	10	1010.5	Partly cloudy	
Stavropol, R.S.F.S.R.	45° 02'	41° 54'	150	1952	Sept.	15	15.0	1.5	75	10	1010.5	Partly cloudy	
Novozybkov, R.S.F.S.R.	52° 32'	31° 56'	150	1952	Sept.	13	15.0	1.5	75	10	1010.5	Partly cloudy	
Meleuz, R.S.F.S.R.	52° 58'	55° 55'	150	1952	Sept.	5	15.0	1.5	75	10	1010.5	Partly cloudy	
Moscow, R.S.F.S.R.	55° 50'	37° 33'	150	1952	Aug.	27	15.0	1.5	75	10	1010.5	Partly cloudy	
Dyurtyuli, R.S.F.S.R.	55° 29'	54° 54'	150	1952	Aug.	10	15.0	1.5	75	10	1010.5	Partly cloudy	
Staraya Chartoriya, Ukrainian S.S.R.	50° 00'	27° 42'	150	1952	Sept.	14	15.0	1.5	75	10	1010.5	Partly cloudy	
Bel-Agach, Kazakh S.S.R.	50° 48'	80° 44'	150	1952	Aug.	26	15.0	1.5	75	10	1010.5	Partly cloudy	
Chakino, R.S.F.S.R.	52° 14'	41° 38'	150	1952	Aug.	30	15.0	1.5	75	10	1010.5	Partly cloudy	
Kazan, R.S.F.S.R.	55° 47'	49° 08'	150	1952	Aug.	12	15.0	1.5	75	10	1010.5	Partly cloudy	
Ust-Kamenogorsk, Kazakh S.S.R.	49° 45'	82° 41'	150	1952	Aug.	28	15.0	1.5	75	10	1010.5	Partly cloudy	
Novo-Urensk, R.S.F.S.R.	54° 32'	52° 48'	150	1952	Aug.	25	15.0	1.5	75	10	1010.5	Partly cloudy	
Bugulma, R.S.F.S.R.	56° 06'	69° 28'	150	1952	Sept.	5	15.0	1.5	75	10	1010.5	Partly cloudy	
Ishim, R.S.F.S.R.	54° 34'	55° 22'	150	1952	Aug.	26	15.0	1.5	75	10	1010.5	Partly cloudy	
Chishmy, R.S.F.S.R.	56° 05'	63° 38'	150	1952	Sept.	3	15.0	1.5	75	10	1010.5	Partly cloudy	
Shadrinsk, R.S.F.S.R.	58° 20'	55° 10'	150	1952	Aug.	18	15.0	1.5	75	10	1010.5	Partly cloudy	
Mendeleyev, R.S.F.S.R.	53° 46'	44° 58'	150	1952	Aug.	25	15.0	1.5	75	10	1010.5	Partly cloudy	
Anuchino, R.S.F.S.R.	58° 22'	51° 35'	150	1952	Aug.	14	15.0	1.5	75	10	1010.5	Partly cloudy	
Falenki, R.S.F.S.R.	53° 20'	83° 48'	150	1952	Aug.	25	15.0	1.5	75	10	1010.5	Partly cloudy	
Barnaul, R.S.F.S.R.	56° 38'	57° 42'	150	1952	Sept.	3	15.0	1.5	75	10	1010.5	Partly cloudy	
Krasnoufimsk, R.S.F.S.R.	55° 46'	43° 54'	150	1952	Sept.	6	15.0	1.5	75	10	1010.5	Partly cloudy	
Surovatikha, R.S.F.S.R.	57° 50'	39° 20'	150	1952	Sept.	29	15.0	1.5	75	10	1010.5	Partly cloudy	
Yaroslavl, R.S.F.S.R.	59° 25'	30° 15'	150	1952	Sept.	10	15.0	1.5	75	10	1010.5	Partly cloudy	
Shelton, R.S.F.S.R.	53° 54'	27° 33'	150	1952	Aug.	25	15.0	1.5	75	10	1010.5	Partly cloudy	
Minsk, White Russian S.S.R.	54° 58'	73° 20'	150	1952	Aug.	24	15.0	1.5	75	10	1010.5	Partly cloudy	
Omsk, R.S.F.S.R.	58° 12'	68° 14'	150	1952	Aug.	21	15.0	1.5	75	10	1010.5	Partly cloudy	
Tobolsk, R.S.F.S.R.	52° 31'	68° 46'	150	1952	Aug.	21	15.0	1.5	75	10	1010.5	Partly cloudy	
Balkashino, Kazakh S.S.R.	53° 00'	37° 23'	150	1952	Sept.	3	15.0	1.5	75	10	1010.5	Partly cloudy	
Shatilov, R.S.F.S.R.	59° 50'	56° 10'	150	1952	Aug.	22	15.0	1.5	75	10	1010.5	Partly cloudy	
Solikam, R.S.F.S.R.	59° 34'	30° 07'	150	1952	Aug.	29	15.0	1.5	75	10	1010.5	Partly cloudy	
Gatchina, R.S.F.S.R.	61° 40'	53° 10'	150	1952	Aug.	27	15.0	1.5	75	10	1010.5	Partly cloudy	
Ust-Kulom, R.S.F.S.R.	59° 40'	34° 15'	150	1952	Aug.	26	15.0	1.5	75	10	1010.5	Partly cloudy	
Tikhvin, R.S.F.S.R.	60° 40'	34° 30'	150	1952	Aug.	29	15.0	1.5	75	10	1010.5	Partly cloudy	
Podporozhye, R.S.F.S.R.	61° 16'	46° 35'	150	1952	Aug.	25	15.0	1.5	75	10	1010.5	Partly cloudy	
Kotlas, R.S.F.S.R.	57° 41'	93° 18'	150	1952	Aug.	26	15.0	1.5	75	10	1010.5	Partly cloudy	
Kazachinskoye, R.S.F.S.R.	56° 26'	84° 58'	150	1952	Sept.	2	15.0	1.5	75	10	1010.5	Partly cloudy	
*Tomsk, R.S.F.S.R.				1952	Aug.	15	15.0	1.5	75	10	1010.5	Partly cloudy	

Source: Based on data as shown in Tables 24 to 32 incl. and additional data of agricultural experiment stations of the Soviet Union between 1898 and 1952, with an over-all average of 5 years for each station. *Indicates stations for which day-degree summations were computed. See Tables 24 to 32 incl.

† Data for local varieties of wheat.

‡ Irrigated.

n.a. - not available.

Figure 18




TABLE 5

AVERAGE DATES AND EXTREMES OF SOWING, HEADING, AND RIPENING OF SPRING WHEAT¹ AT AGRICULTURAL EXPERIMENT STATIONS IN THE SOVIET UNION

(Arranged from earliest to latest average ripening dates)

LOCATION	LAT. (N)	LONG. (E)	DATE SOWN			DATE HEADED			DATE RIPE		
			AVERAGE	EXTREMES	AVERAGE	AVERAGE	EXTREMES	AVERAGE	AVERAGE	EXTREMES	EXTREMES
*Samarkand, Uzbek S.S.R.	39° 39'	66° 57'	Mar. 6	Feb. 20—Mar. 20	n.a.	June 6	—	June 27	June 18—July 1		
*Kherson, Ukrainian S.S.R.	46° 38'	32° 37'	Mar. 29	Mar. 15—Apr. 15	n.a.	June 6	—	June 27	June 27—July 14		
Zhdanov, Ukrainian S.S.R.	47° 06'	37° 35'	Apr. 10	Apr. 5—Apr. 14	June 7	June 7	May 28—June 15	July 4	July 1—July 19		
*Krasnodar, R.S.F.S.R.	45° 02'	38° 56'	Mar. 8	Feb. 5—Apr. 9	n.a.	n.a.	—	July 9	June 29—July 18		
Andizhan, Uzbek S.S.R.	40° 45'	72° 22'	Mar. 22	n.a.	n.a.	n.a.	—	July 10	n.a.		
Rostov-Na-Donu, R.S.F.S.R.	47° 13'	39° 43'	Apr. 5	Mar. 28—Apr. 14	June 8	June 8	June 5—June 11	July 11	July 7—July 13		
Yevsk, R.S.F.S.R.	46° 40'	38° 16'	Apr. 9	Mar. 27—Apr. 21	June 12	June 12	June 4—June 19	July 12	July 4—July 20		
Persyanovka, R.S.F.S.R.	47° 32'	40° 07'	Apr. 13	Apr. 10—Apr. 16	...do...	...do...	June 9—June 14	...do...	July 9—July 16		
Kirovograd, Ukrainian S.S.R.	48° 31'	32° 17'	Apr. 5	Apr. 1—Apr. 9	June 11	June 11	June 10—June 19	...do...	July 9—July 19		
Milyutinskaya, R.S.F.S.R.	48° 37'	41° 41'	Apr. 17	n.a.	June 8	June 8	n.a.	July 13	n.a.		
Starobelsk, Ukrainian S.S.R.	49° 16'	38° 56'	Apr. 24	Apr. 14—May 5	June 12	June 12	June 5—June 25	...do...	July 8—July 21		
Mirgorod, Ukrainian S.S.R.	49° 58'	33° 37'	n.a.	n.a.	June 15	June 15	n.a.	...do...	July 5—July 19		
Khasavyurt, R.S.F.S.R.	43° 24'	46° 10'	Apr. 14	n.a.	June 17	June 17	June 11—June 23	July 14	July 10—July 18		
Voroshilovgrad, Ukrainian S.S.R.	48° 35'	39° 20'	Apr. 5	n.a.	June 13	June 13	June 5—June 15	...do...	July 8—July 18		
Khorol, Ukrainian S.S.R.	49° 48'	33° 15'	Apr. 15	Apr. 10—Apr. 19	June 15	June 15	June 10—June 19	...do...	July 9—July 19		
Sinelnikovo, Ukrainian S.S.R.	48° 21'	35° 26'	n.a.	n.a.	...do...	...do...	n.a.	July 15	July 8—July 19		
Drabov, Ukrainian S.S.R.	49° 57'	32° 19'	Apr. 15	Apr. 10—Apr. 20	...do...	...do...	June 10—June 20	July 16	July 6—July 20		
Adzamka, Ukrainian S.S.R.	48° 34'	32° 24'	Apr. 14	Apr. 13—Apr. 15	...do...	...do...	...do...	July 17	July 10—July 30		
Belaya Tserkov, Ukrainian S.S.R.	49° 47'	30° 09'	Apr. 21	Apr. 17—Apr. 25	June 18	June 18	June 10—June 29	...do...	July 6—July 20		
Dessa, Ukrainian S.S.R.	46° 29'	30° 44'	Apr. 9	Mar. 25—May 2	June 14	June 14	June 7—June 26	July 18	July 11—July 27		
Nemerchi, Ukrainian S.S.R.	48° 40'	27° 45'	n.a.	—	June 12	June 12	June 5—June 15	July 18	July 15—July 25		
Gumy, Ukrainian S.S.R.	50° 53'	34° 43'	Apr. 20	Apr. 10—Apr. 30	June 18	June 18	June 15—June 25	July 21	July 18—July 25		
Saratov, R.S.F.S.R.	51° 32'	46° 01'	Apr. 23	Apr. 8—May 6	...do...	...do...	June 2—July 1	...do...	June 29—Aug. 2		
Lamenets-Podolsk, Ukrainian S.S.R.	48° 40'	26° 34'	n.a.	—	n.a.	n.a.	—	July 22	July 10—July 29		
Trasnyy Kut, R.S.F.S.R.	50° 58'	46° 30'	Apr. 21	Apr. 12—May 6	June 15	June 15	May 30—June 29	...do...	July 4—Aug. 3		
Iershov, R.S.F.S.R.	51° 20'	48° 17'	May 2	Apr. 29—May 4	June 20	June 20	June 16—June 24	...do...	July 17—July 27		
Ioty, Moldavian S.S.R.	47° 57'	29° 10'	Apr. 2	Mar. 20—Apr. 24	June 17	June 17	June 7—June 30	July 24	July 17—Aug. 6		
Olyanka, R.S.F.S.R.	51° 30'	50° 05'	Apr. 30	Apr. 25—May 4	June 22	June 22	June 18—June 25	...do...	July 22—July 27		
harkov, Ukrainian S.S.R.	49° 58'	36° 09'	Apr. 15	n.a.	June 18	June 18	June 11—June 30	July 25	July 3—Aug. 4		
emir, Kazakh S.S.R.	49° 06'	57° 10'	Apr. 30	Apr. 18—May 11	n.a.	n.a.	—	...do...	July 15—Aug. 14		
ralsk, R.S.F.S.R.	51° 12'	51° 21'	May 3	Apr. 24—May 12	June 24	June 24	June 17—July 2	July 26	July 14—Aug. 7		
Itava, Ukrainian S.S.R.	49° 35'	34° 34'	Apr. 27	Apr. 14—May 5	June 20	June 20	June 14—June 25	July 27	July 19—July 30		
dashov, R.S.F.S.R.	51° 33'	43° 09'	Apr. 30	n.a.	June 22	June 22	June 21—June 23	July 28	July 25—Aug. 1		
avropol, R.S.F.S.R.	45° 02'	41° 54'	Apr. 19	Mar. 11—May 18	June 23	June 23	June 13—July 8	July 30	July 20—Aug. 10		
zenchuk, R.S.F.S.R.	53° 00'	49° 28'	Apr. 26	Apr. 21—Apr. 29	June 27	June 27	June 20—July 2	...do...	July 20—Aug. 8		
Glyshchev, R.S.F.S.R.	53° 12'	50° 09'	Apr. 19	Apr. 11—Apr. 27	...do...	...do...	June 26—June 28	...do...	July 25—Aug. 3		
ronozh, R.S.F.S.R.	51° 40'	39° 13'	Apr. 16	Apr. 2—Apr. 30	June 20	June 20	June 17—June 24	July 31	July 29—Aug. 2		
ishmy, R.S.F.S.R.	54° 34'	55° 23'	Apr. 28	Apr. 10—May 13	June 30	June 30	June 14—July 12	Aug. 2	July 13—Sept. 10		
nenka, R.S.F.S.R.	53° 48'	83° 34'	May 6	Apr. 29—May 14	n.a.	n.a.	—	Aug. 4	Aug. 2—Aug. 7		
ndiro, R.S.F.S.R.	54° 05'	91° 07'	May 5	Apr. 29—May 14	n.a.	n.a.	—	Aug. 4	Aug. 2—Aug. 7		

~~SECRET~~

Table 4

Approximate Seeding and Harvesting Dates for Oats
in Specified Districts of the USSR 5/

<u>Specified District a/</u>	<u>Seeding Season</u>		<u>Harvesting Season</u>	
	<u>Beginning</u>	<u>Ending</u>	<u>Beginning</u>	<u>Ending</u>
<u>European USSR</u>				
Chief Wheat Surplus Area				
South Ukraine (III)	15 Mar	20 Apr	15 Jul	25 Jul
Lower Don-North Caucasus (IV)	20 Mar	15 May	15 Jul	30 Jul
Lower Volga (VI)	1 Apr	5 May	20 Jul	5 Aug
Middle Volga (VI)	20 Apr	15 May	10 Aug	30 Aug
Bashkir ASSR (VIII)	10 Apr	20 May	20 Aug	10 Sep
Chief Rye Surplus Area				
West and North Ukraine (III)	20 Mar	1 May	20 Jul	10 Aug
Black Soil Belt (VII)	25 Mar	15 May	30 Jul	15 Aug
Northern Deficit Area				
Belorussian SSR (IIb)	10 Apr	30 May	15 Aug	5 Oct
Central Industrial Concentration (VII)	10 Apr	20 May	20 Aug	10 Sep
Tatar ASSR (VI)	10 Apr	20 May	20 Aug	20 Sep
Baltic States and North-Western (IIa and Ia)	15 Apr	10 Jun	20 Aug	10 Sep
Northern European USSR (Ib)	10 Apr	10 Jun	1 Sep	20 Sep
<u>Asiatic USSR</u>				
Spring Wheat Surplus Belt				
Ural East of the Ural Mountains (VIII and IX)	10 Apr	10 Jun	20 Aug	15 Sep
West Siberia, Southern Part (IX)	10 Apr	10 Jun	20 Aug	15 Sep
Kazakh SSR, Northern Part (Xa)	10 Apr	10 Jun	25 Aug	N.A.
Central Asia				
Tajik SSR (Xb)	b/	b/	b/	b/
Kirgiz SSR (Xb)	1 Apr	20 May	30 Jul	N.A.
Transcaucasus (V)	b/	b/	b/	b/

a. The numerals in parentheses refer to the Economic Regions shown on the map following p. 18.

b. The acreage seeded, if any, is negligible.